

REFERENCES

- Aikaite-Stanaitiene, J., Grigiškis, S., Levišauskas, D., Čipinyte, V., Baškys, E., & Kačkyte, V. (2010). Development of fatty waste composting technology using bacterial preparation with lipolytic activity. *Journal of Environmental Engineering and Landscape Management*, 18(4), 296-305.
- Al-Darbi, M. M., Saeed, N. O., Islam, M. R., & Lee, K. (2005). Biodegradation of Natural Oils in Seawater. *Energy Sources*, 27(1-2), 19–34. doi,10.1080/00908310490448073.
- Alexandre, V. M. F., Valente, A. M., Cammarota, M. C., & Freire, D. M. (2011). Performance of anaerobic bioreactor treating fish-processing plant wastewater pre-hydrolyzed with a solid enzyme pool. *Renewable energy*, 36(12), 3439-3444.
- Alexandria, VA. Cipnyte, V., Grigiškis, S., Baškys, E. (2009). Selection of fat-degrading microorganisms for the treatment of lipid-contaminated environment. *BIOLOGIJA*, 55(3–4), 84–92.
- Alves, M. M., Vieira, J. M., Pereira, R. A., Pereira, M. A., & Mota, M. (2001). Effects of lipids and oleic acid on biomass development in anaerobic fixed-bed reactors. Part II, Oleic acid toxicity and biodegradability. *Water Research*, 35(1), 264-270.
- Bligh, E. G., & Dyer, W. J. (1959). A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology*, 37(8), 911-917.
- Bustillo-Lecompte, C. F., Mehrvar, M., & Quiñones-Bolaños, E. (2014). Cost-effectiveness analysis of TOC removal from slaughterhouse wastewater using combined anaerobic-aerobic and UV/H₂O₂ processes. *Journal of Environmental Management*, 134, 145-152.
- Caixeta, C. E., Cammarota, M. C., & Xavier, A. M. (2002). Slaughterhouse wastewater treatment, evaluation of a new three-phase separation system in a UASB reactor. *Bioresource Technology*, 81(1), 61-69.
- Cammarota, M. C., and D. M. G. Freire. (2006). A review on hydrolytic enzymes in the treatment of wastewater with high oil and grease content. *Bioresource Technology* 97(17), 2195-2210.
- Cammarota, M.C., Freire, D.M.G., Sant'Anna Jr., G.L., Russo, C., Freire, D.D.C., Castilho, L.R. (2003). Production process and composition of an enzymatic preparation and its use for the treatment of domestic and industrial effluents of high fat, protein and/or carbohydrate content. Patent PCT/BR01/00124, New Zealand.
- Cammarota, M.C., Teixeira, G.A., Freire, D.M.G. (2001). Enzymatic prehydrolysis and anaerobic degradation of wastewaters with high oil contents. *Biotechnology Letters*, 23, 1591–1595.
- Chan, H. (2010). Removal and recycling of pollutants from Hong Kong restaurant wastewaters. *Bioresource Technology*, 101(17), 6859-67.

- de Castro, P. F., Moreira, N. C., Esperan  a, M. N., de Oliveira, L. M., Badino, A. C., Tavano, O. L., & Hirata, D. B. (2016). High Lipase Production from *Geotrichum candidum* in Reduced Time using Cottonseed Oil, Optimization, Easy Purification and Specificity Characterization. *Journal of Chemical Engineering Research Updates*, 3, 60-69.
- Choe, E., Min, D.,B. (2007). Chemistry of deep-fat frying oils. *Journal of food science*. Jun 1;72(5), 77-86.
- Christie, W.W. (1993). Preparation of ester derivatives of fatty acids for chromatographic analysis. *Advances in Lipid Methodology*, 2(69), 111.
- Chu, W. & Hsu, W. (1999). Pollution source identification and waste loading reduction at Chinese fast food restaurants. *Environment international*, 25(1), 97-107.
- Chung, G., Downing, L., Kabouris, J., & Shea, T. (2010). Direct Addition of High-Strength Organic Waste to Municipal Wastewater Anaerobic Digesters. *Water Environment Federation, Tech. Alexandria, VA*.
- Costa, T. M., Hermann, K. L., Garcia-Roman, M., Valle, R. D. C. S. C., & Tavares, L. B. B. (2017). Lipase Production by *Aspergillus niger* Grown in Different Agro-Industrial Wastes by Solid-State Fermentation. *Brazilian Journal of Chemical Engineering*, 34(2), 419-427.
- Cvengroš J., Cvengrošov   Z. (2004). Used frying oils and fats and their utilization in the production of methyl esters of higher fatty acids. *Biomass and Bioenergy*. 227(2), 173-81.
- Danalewich, J. R., Papagiannis, T. G., Belyea, R. L., Tumbleson, M. E., & Raskin, L. (1998). Characterization of dairy waste streams, current treatment practices, and potential for biological nutrient removal. *Water Research*, 32(12), 3555-3568.
- Dassey, A.J., and C.S. Theegala. (2012). Evaluating coagulation pretreatment on poultry processing wastewater for dissolved air flotation. *Journal of Environmental Science and Health, Part A* 47(13), 2069-2076.
- de Moraes J  nior, W. G., Kamimura, E. S., Ribeiro, E. J., Pessela, B. C., Cardoso, V. L., & de Resende, M. M. (2016). Optimization of the production and characterization of lipase from *Candida rugosa* and *Geotrichum candidum* in soybean molasses by submerged fermentation. *Protein expression and purification*, 123, 26-34.
- Del Nery, V., De Nardi, I. R., Damianovic, M. H. R. Z., Pozzi, E., Amorim, A. K. B., & Zaiat, M. (2007). Long-term operating performance of a poultry slaughterhouse wastewater treatment plant. *Resources, conservation and recycling*, 50(1), 102-114.
- El-Bestawy, E., El-Masry, M. H., & Nawal, E. (2005). The potentiality of free Gram-negative bacteria for removing oil and grease from contaminated industrial effluents. *World Journal of Microbiology and Biotechnology*, 21(6-7), 815-822.
- Falony G., Armas J.C., Mendoza J.C, Hern  ndez J.L. (2006). Production of extracellular lipase from *Aspergillus niger* by solid-state fermentation. *Food Technology and Biotechnology*. 44(2), 235-240.

- Federation, W. E., & American Public Health Association. (2012). *Standard Methods for the Examination of Water and Wastewater*. 22nd Edition, American Public Health Association (APHA), Washington, DC, USA.
- Gaur, R.S., Cai, L., Tuovinen, O.H., Mancl, K.M. (2010). Pretreatment of turkey fat-containing wastewater in coarse sand and gravel/coarse sand bioreactors. *Bioresource Technology*, 101, 1106–1110.
- Gombert, A. K., Pinto, A. L., Castilho, L. R., & Freire, D. M. (1999). Lipase production by *Penicillium restrictum* in solid-state fermentation using babassu oil cake as substrate. *Process Biochemistry*, 35(1), 85-90.
- Hanaki, K., Matsuo, T., & Nagase, M. (1981). Mechanism of inhibition caused by long-chain fatty acids in anaerobic digestion process. *Biotechnology and Bioengineering*, 23(7), 1591-1610.
- Hee-Yeon, C. H. O., Bancerz, R., Ginalska, G., Leonowicz, A., Nam-Seok, C. H. O., & Shoji, O. H. G. A. (2007). Culture conditions of psychrotrophic fungus, *Penicillium chrysogenum* and its lipase characteristics. *Journal of Faculty of Agriculture, Kyushu University*, 52(2), 281-286.
- Hwu, C. S., & Lettinga, G. (1997). Acute toxicity of oleate to acetate-utilizing methanogens in mesophilic and thermophilic anaerobic sludges. *Enzyme and Microbial Technology*, 21(4), 297-301.
- Ivanov, V. N., Stabnikova, E. V., Stabnikov, V. P., Kim, I. S., & Zubair, A. (2002). Effects of iron compounds on the treatment of fat-containing wastewaters. *Applied Biochemistry and Microbiology*, 38(3), 255-258.
- Janczukowicz, W., Zieliński, M., & Dębowksi, M. (2008). Biodegradability evaluation of dairy effluents originated in selected sections of dairy production. *Bioresource Technology*, 99(10), 4199-4205.
- Jeganathan, J., Bassi, A., & Nakhla, G. (2006). Pre-treatment of high oil and grease pet food industrial wastewaters using immobilized lipase hydrolyzation. *Journal of Hazardous Materials*, 137(1), 121–128. Jung, F., Cammarota, M. C., & Freire, D. M. G. (2002). Impact of enzymatic pre-hydrolysis on batch activated sludge systems dealing with oily wastewaters. *Biotechnology Letters*, 24(21), 1797-1802.
- Kang, J. X., Lu, L., Zhan, W., Li, B., Li, D. S., Ren, Y. Z., & Liu, D. Q. (2011). Photocatalytic pretreatment of oily wastewater from the restaurant by a vacuum ultraviolet/TiO₂ system. *Journal of hazardous materials*, 186(1), 849-854.
- Keenan, D., & Sabelnikov, A. (2000). Biological augmentation eliminates grease and oil in bakery wastewater. *Water Environment Research*, 72(2), 141-146.
- Kiepper, B., Governo, J., & Zacharias, H. (2011). Characterization of the generation, handling and treatment of spent fat, oil, and grease (FOG) from Georgia's food service industry. College of Agricultural & Environmental Sciences, Department of Biological & Agricultural Engineering, Engineering Outreach Program, University of Georgia.

- Kim, S. H., Han, S. K., & Shin, H. S. (2004). Two-phase anaerobic treatment system for fat-containing wastewater. *Journal of Chemical Technology and Biotechnology*, 79(1), 63-71.
- Kim, S.H., Shin, H.S., (2010). Enhanced lipid degradation in an upflow anaerobic sludge blanket reactor by integration with an acidogenic reactor. *Water Environ. Res.* 82, 267–272.
- Kobayashi, T., Xu, K.Q., Li, Y.Y., Inamori, Y. (2012). Effect of sludge recirculation on characteristics of hydrogen production in a two-stage hydrogen–methane fermentation process treating food wastes. *International Journal of Hydrogen Energy*, 37, 5602–5611.
- Koster, I. W. (1987). Abatement of long-chain fatty acid inhibition of methanogenesis by calcium addition. *Biological Wastes*, 22(4), 295-301.
- Kotogán, A., Németh, B., Vágvölgyi, C., Papp, T., & Takó, M. (2014). Screening for extracellular lipase enzymes with transesterification capacity in Mucoromycotina strains. *Food Technology and Biotechnology*, 52(1), 73-82.
- Kowalski, R. (2007). GC analysis of changes in the fatty acid composition of sunflower and olive oils heated with quercetin, caffeic acid, protocatechuic acid, and butylated hydroxyanisole. *Acta Chromatographica*, 18, 15.
- Kumar, A., & Kanwar, S. S. (2012). Lipase production in solid-state fermentation (SSF), recent developments and biotechnological applications. *Dynamic Biochemistry, Process Biotechnology and Molecular Biology*, 6(1), 13-27.
- Kumar, S., Mathur, A., Singh, V., Nandy, S., Khare, S. K., & Negi, S. (2012). Bioremediation of waste cooking oil using a novel lipase produced by *Penicillium chrysogenum*SNP5 grown in solid medium containing waste grease. *Bioresource Technology*, 120, 300-304.
- Kuo, W. C., & Cheng, K. Y. (2007). Use of respirometer in evaluation of process and toxicity of thermophilic anaerobic digestion for treating kitchen waste. *Bioresource Technology*, 98(9), 1805-1811.
- Lam, M. K., Lee, K. T., & Mohamed, A. R. (2010). Homogeneous, heterogeneous and enzymatic catalysis for transesterification of high free fatty acid oil (waste cooking oil) to biodiesel, a review. *Biotechnology Advances*, 28(4), 500-518.
- Leal, M.C.M.R., Cammarota, M.C.M., Freire, D.M.G., Sant'Anna Jr., G.L. (2002). Hydrolytic enzymes as coadjuvants in the anaerobic treatment of dairy wastewaters. *Braz. Journal of Chemical Engineering*, 19, 175–180.
- Li, Z., Wrenn, B.A., Venosa, A.D., 2005. Effect of iron on the sensitivity of hydrogen, acetate, and butyrate metabolism to inhibition by long-chain fatty acids in vegetable-oil-enriched freshwater sediments. *Water Research*. 39, 3109-3119.
- Lin, Y., Wang, D., Wu, S. and Wang, C. (2009). Alkali pretreatment enhances biogas production in the anaerobic digestion of pulp and paper sludge. *Journal of Hazardous Materials*, 170(1), 366-373.

- Long, J. H., Aziz, T. N., Francis, L., & Ducoste, J. J. (2012). Anaerobic co-digestion of fat, oil, and grease (FOG), a review of gas production and process limitations. *Process Safety and Environmental Protection*, 90(3), 231-245.
- Mahadik, N. D., Puntambekar, U. S., Bastawde, K. B., Khire, J. M., & Gokhale, D. V. (2002). Production of acidic lipase by *Aspergillus niger* in solid state fermentation. *Process Biochemistry*, 38(5), 715-721.
- Mahmoud, G. A. E., Koutb, M. M., Morsy, F. M., & Bagy, M. M. (2015). Characterization of lipase enzyme produced by hydrocarbons utilizing fungus *Aspergillus terreus*. *European Journal of Biological Research*, 5(3), 70-77.
- Masse, L., Kennedy, K. J., & Chou, S. (2001). Testing of alkaline and enzymatic hydrolysis pretreatments for fat particles in slaughterhouse wastewater. *Bioresource Technology*, 77(2), 145-155.
- Mendes, A.A., Castro, H.F., Pereira, E.B., Furigo Jr., A. (2005). Application of lipases for wastewater treatment containing high levels of lipids. *Química Nova*, 28, 296–305.
- Metcalf E, Eddy E. *Wastewater Engineering, Treatment and Reuse*. McGrawHill. Inc., New York. 2003.
- Mouneimne, A. H., Carrere, H., Bernet, N., & Delgenes, J. P. (2004). Effect of the addition of bentonite on the anaerobic biodegradability of solid fatty wastes. *Environmental Technology*, 25(4), 459-469.
- Mukhtar, H., Hanif, M. A. D. I. H. A., Rehman, A. U., Nawaz, A., & Haq, I. U. (2015). Studies on the Lipase Production by *Aspergillus niger* through Solid State Fermentation. *Pakistan Journal of Botany*, 47, 351-354.
- Nanayakkara, C.M., Witharana, A. (2015). Bioremediation of Oil Contaminated Soil and Water, In situ and Ex situ Strategies for Feasibility Assessment in Singh S. & Srivastava K. (Eds.) *Handbook of Research on Uncovering New Methods for Ecosystem Management through Bioremediation*. IGI Global. USA. pp 222-254.
- Nordell, E., Hansson, A. B., & Karlsson, M. (2013). Zeolites relieves inhibitory stress from high concentrations of long chain fatty acids. *Waste management*, 33(12), 2659-2663.
- Omar, T.A., Salimon, J. (2013). Validation and application of a gas chromatographic method for determining fatty acids and trans fats in some bakery products, *Journal of Taibah University for Science*. <http://dx.doi.org/10.1016/j.jtusci.2013.04.002>.
- Palatsi, J., Lauren, M., Andrés, M.V., Flotats, X., Nielsen, H.B., Angelidaki, I., (2009). Strategies for recovering inhibition caused by long chain fatty acids on anaerobic thermophilic biogas reactors. *Bioresource Technology*. 100, 4588e4596.
- Palatsi, J., Lauren, M., Andrés, M. V., Flotats, X., Nielsen, H. B., & Angelidaki, I. (2009). Strategies for recovering inhibition caused by long chain fatty acids on anaerobic thermophilic biogas reactors. *Bioresource Technology*, 100(20), 4588-4596.
- Parihar, D. K. (2012). Production of lipase utilizing linseed oilcake as fermentation substrate. *International Journal of Science and Technoogy*, 1(3), 135-143.

- Pereira, M.A., Sousa, D.Z., Mota, M., Alves, M.M., 2004. Mineralization of LCFA associated with anaerobic sludge kinetics, enhancement of methanogenic activity, and effect of VFA. *Biotechnol. Bioeng.* 88, 502-511.
- Pintor, A. M., Vilar, V. J., Botelho, C. M., & Boaventura, R. A. (2016). Oil and grease removal from wastewaters, Sorption treatment as an alternative to state-of-the-art technologies,A critical review. *Chemical Engineering Journal*, 297, 229-255.
- Raimbault, M., & Alazard, D. (1980). Culture method to study fungal growth in solid fermentation. *Applied Microbiology and Biotechnology*, 9(3), 199-209.
- Rashid, N., Imanaka, T. (2008). Efficient degradation of grease using microorganisms. *Journal of Chemical Society of Pakistan*, 30(4), 612-617.
- Rashid, M.T., Voroney, R.P. (2004). Land application of oily food waste and corn production on amended soils. *Agronomy Journal*, 96(4), 997–1004.
- Ratnayake, N. and Miththapala, S. (2011). Study on sustainable consumption practices in Sri Lanka hotel sector. Proceedings of Civil Engineering Research for Industry. Department of Civil Engineering, University of Moratuwa, Moratuwa.
- Rosa, D. R., Duarte, I. C., Saavedra, N. K., Varesche, M. B., Zaiat, M., Cammarota, M. C., & Freire, D. M. (2009). Performance and molecular evaluation of an anaerobic system with suspended biomass for treating wastewater with high fat content after enzymatic hydrolysis. *Bioresource Technology*, 100(24), 6170-6176.
- Santis-Navarro, A., Gea, T., Barrena, R., & Sánchez, A. (2011). Production of lipases by solid state fermentation using vegetable oil-refining wastes. *Bioresource Technology*, 102(21), 10080-10084.
- Sarkar, Baisali, P. P. Chakrabarti, A. Vijaykumar, and Vijay Kale. (2006).Wastewater treatment in dairy industries-possibility of reuse. *Desalination* 195, no. 1-3, 141-152.
- Schmidt, Curtis J. (1974). Wastewater characterization for the specialty food industry.
- Sharma, S. (2012). Bioremediation, features, strategies and applications. *Asian Journal of Pharmacy and Life Science*, ISSN2231, 4423.
- Shin, H.S., Kim, S.H., Lee, C.Y., Nam, S.Y. (2003). Inhibitory effects of long-chain fatty acids on VFA degradation and β -oxidation. *Water Science and Technology*.47(10),139-46.
- Silvestre, G., A. Rodríguez-Abalde, B. Fernández, X. Flotats, and A. Bonmatí. (2011). Biomass adaptation over anaerobic co-digestion of sewage sludge and trapped grease waste. *Bioresource Technology* 102(13), 6830-6836.
- Tocchi, C., Federici, E., Fidati, L., Manzi, R., Vinciguerra, V., & Petruccioli, M. (2012). Aerobic treatment of dairy wastewater in an industrial three-reactor plant, Effect of aeration regime on performances and on protozoan and bacterial communities. *WaterResearch*, 46(10), 3334-3344.

- Treichel, H., Sbardelotto, M., Venturin, B., Agnol, A. D., Mulinari, J., Golunski, S. M., & Vargas, G. D. (2017). Lipase production from a newly isolated *Aspergillus niger* by solid state fermentation using canola cake as substrate. *Current Biotechnology*, 6(4), 295-300.
- Ülker, S., Öznel, A., Colak, A., & Karaoğlu, Ş. A. (2011). Isolation, production, and characterization of an extracellular lipase from *Trichoderma harzianum* isolated from soil. *Turkish Journal of Biology*, 35(5), 543-550.
- Valladão, A. B. G., Freire, D. M. G., & Cammarota, M. C. (2007). Enzymatic pre-hydrolysis applied to the anaerobic treatment of effluents from poultry slaughterhouses. *International Biodeterioration & Biodegradation*, 60(4), 219-225.
- Wakelin, N. G., & Forster, C. F. (1998). The aerobic treatment of grease-containing fast food restaurant wastewaters. *Process Safety and Environmental Protection*, 76(1), 55-61.
- Weng, C.N., Jeris, J.S., (1976). Biochemical mechanisms in the methane fermentation of glutamic and oleic acids. *Water Research*. 10, 9-18.
- Wu, L. J., Kobayashi, T., Kuramochi, H., Li, Y. Y., & Xu, K. Q. (2015). Recovery strategies of inhibition for mesophilic anaerobic sludge treating the de-oiled grease trap waste. *International Biodeterioration & Biodegradation*, 104, 315-323.