

REFERENCES

- [1] M.A. Borowitzka, High-value products from microalgae—their development and commercialisation, *J. Appl. Phycol.* 25 (2013) 743–756.
<https://doi.org/10.1007/s10811-013-9983-9>.
- [2] M. Koller, A. Muhr, G. Braunegg, Microalgae as versatile cellular factories for valued products, *Algal Res.* 6 (2014) 52–63.
<https://doi.org/10.1016/j.algal.2014.09.002>.
- [3] C.-Y. Chen, X.-Q. Zhao, H.-W. Yen, S.-H. Ho, C.-L. Cheng, D.-J. Lee, F.-W. Bai, J.-S. Chang, Microalgae-based carbohydrates for biofuel production, *Biochem. Eng. J.* 78 (2013) 1–10. <https://doi.org/10.1016/j.bej.2013.03.006>.
- [4] Y. Chisti, Biodiesel from microalgae, *Biotechnol. Adv.* 25 (2007) 294–306.
<https://doi.org/10.1016/j.biotechadv.2007.02.001>.
- [5] I. Hariskos, C. Posten, Biorefinery of microalgae - opportunities and constraints for different production scenarios, *Biotechnol. J.* 9 (2014) 739–752.
<https://doi.org/10.1002/biot.201300142>.
- [6] K.W. Chew, J.Y. Yap, P.L. Show, N.H. Suan, J.C. Juan, T.C. Ling, D.-J. Lee, J.-S. Chang, Microalgae biorefinery: High value products perspectives, *Bioresour. Technol.* 229 (2017) 53–62. <https://doi.org/10.1016/j.biortech.2017.01.006>.
- [7] L.M. Schüler, P.S.C. Schulze, H. Pereira, L. Barreira, R. León, J. Varela, Trends and strategies to enhance triacylglycerols and high-value compounds in microalgae, *Algal Res.* 25 (2017) 263–273.
<https://doi.org/10.1016/j.algal.2017.05.025>.
- [8] Z. Sun, X. Wang, J. Liu, Screening of Isochrysis strains for simultaneous production of docosahexaenoic acid and fucoxanthin, *Algal Res.* 41 (2019) 101545. <https://doi.org/10.1016/j.algal.2019.101545>.
- [9] D.D. McClure, A. Luiz, B. Gerber, G.W. Barton, J.M. Kavanagh, An

- investigation into the effect of culture conditions on fucoxanthin production using the marine microalgae *Phaeodactylum tricornutum*, *Algal Res.* 29 (2018) 41–48. <https://doi.org/10.1016/j.algal.2017.11.015>.
- [10] M. Petrushkina, E. Gusev, B. Sorokin, N. Zotko, A. Mamaeva, A. Filimonova, M. Kulikovskiy, Y. Maltsev, I. Yampolsky, E. Guglya, V. Vinokurov, Z. Namsaraev, D. Kuzmin, Fucoxanthin production by heterokont microalgae, *Algal Res.* 24 (2017) 387–393. <https://doi.org/10.1016/j.algal.2017.03.016>.
- [11] S.M. Kim, S.-W. Kang, O.-N. Kwon, D. Chung, C.-H. Pan, Fucoxanthin as a major carotenoid in *Isochrysis aff. galbana*: Characterization of extraction for commercial application, *J. Korean Soc. Appl. Biol. Chem.* 55 (2012) 477–483. <https://doi.org/10.1007/s13765-012-2108-3>.
- [12] M.-N. Woo, S.-M. Jeon, H.-J. Kim, M.-K. Lee, S.-K. Shin, Y.C. Shin, Y.-B. Park, M.-S. Choi, Fucoxanthin supplementation improves plasma and hepatic lipid metabolism and blood glucose concentration in high-fat fed C57BL/6N mice, *Chem. Biol. Interact.* 186 (2010) 316–322. <https://doi.org/10.1016/j.cbi.2010.05.006>.
- [13] A. Fung, N. Hamid, J. Lu, Fucoxanthin content and antioxidant properties of *Undaria pinnatifida*, *Food Chem.* 136 (2013) 1055–1062. <https://doi.org/10.1016/j.foodchem.2012.09.024>.
- [14] J. Peng, J.-P. Yuan, C.-F. Wu, J.-H. Wang, Fucoxanthin, a Marine Carotenoid Present in Brown Seaweeds and Diatoms: Metabolism and Bioactivities Relevant to Human Health, *Mar. Drugs.* 9 (2011) 1806–1828. <https://doi.org/10.3390/md9101806>.
- [15] S. Xia, K. Wang, L. Wan, A. Li, Q. Hu, C. Zhang, Production, Characterization, and Antioxidant Activity of Fucoxanthin from the Marine Diatom *Odontella aurita*, *Mar. Drugs.* 11 (2013) 2667–2681. <https://doi.org/10.3390/md11072667>.
- [16] F. Gao, I. Teles (Cabanelas, ITD), R.H. Wijffels, M.J. Barbosa, Process

- optimization of fucoxanthin production with *Tisochrysis lutea*, *Bioresour. Technol.* 315 (2020) 123894. <https://doi.org/10.1016/j.biortech.2020.123894>.
- [17] M.L. Hamilton, R.P. Haslam, J.A. Napier, O. Sayanova, Metabolic engineering of *Phaeodactylum tricornutum* for the enhanced accumulation of omega-3 long chain polyunsaturated fatty acids, *Metab. Eng.* 22 (2014) 3–9.
<https://doi.org/10.1016/j.ymben.2013.12.003>.
- [18] J. Liu, M. Sommerfeld, Q. Hu, Screening and characterization of Isochrysis strains and optimization of culture conditions for docosahexaenoic acid production, *Appl. Microbiol. Biotechnol.* 97 (2013) 4785–4798.
<https://doi.org/10.1007/s00253-013-4749-5>.
- [19] E. Ryckebosch, C. Bruneel, K. Muylaert, I. Fouber, Microalgae as an alternative source of omega-3 long chain polyunsaturated fatty acids, *Lipid Technol.* 24 (2012) 128–130. <https://doi.org/10.1002/lite.201200197>.
- [20] E. Ryckebosch, C. Bruneel, R. Termote-Verhalle, K. Goiris, K. Muylaert, I. Fouber, Nutritional evaluation of microalgae oils rich in omega-3 long chain polyunsaturated fatty acids as an alternative for fish oil, *Food Chem.* 160 (2014) 393–400. <https://doi.org/10.1016/j.foodchem.2014.03.087>.
- [21] V. Patil, T. Källqvist, E. Olsen, G. Vogt, H.R. Gislerød, Fatty acid composition of 12 microalgae for possible use in aquaculture feed, *Aquac. Int.* 15 (2007) 1–9.
<https://doi.org/10.1007/s10499-006-9060-3>.
- [22] M.N. Rammuni, T.U. Ariyadasa, P.H.V. Nimarshana, R.A. Attalage, Comparative assessment on the extraction of carotenoids from microalgal sources: Astaxanthin from *H. pluvialis* and β-carotene from *D. salina*, *Food Chem.* 277 (2019) 128–134. <https://doi.org/10.1016/j.foodchem.2018.10.066>.
- [23] M.M.A. Aziz, B. Abdullah, Z. Shokravi, S.A.F. Syed Muhammad, S. Ismail, K.A. Kassim, A.N. Mahmood, H. Shokravi, Two-Stage Cultivation Strategy for Simultaneous Increases in Growth Rate and Lipid Content of Microalgae: A

- Review, Renew. Sustain. Energy Rev. (2019).
- [24] H. Wang, Y. Zhang, L. Chen, W. Cheng, T. Liu, Combined production of fucoxanthin and EPA from two diatom strains *Phaeodactylum tricornutum* and *Cylindrotheca fusiformis* cultures, *Bioprocess Biosyst. Eng.* 41 (2018) 1061–1071. <https://doi.org/10.1007/s00449-018-1935-y>.
 - [25] B. Gao, A. Chen, W. Zhang, A. Li, C. Zhang, Co-production of lipids, eicosapentaenoic acid, fucoxanthin, and chrysolaminarin by *Phaeodactylum tricornutum* cultured in a flat-plate photobioreactor under varying nitrogen conditions, *J. Ocean Univ. China.* 16 (2017) 916–924.
<https://doi.org/10.1007/s11802-017-3174-2>.
 - [26] W. Zhang, F. Wang, B. Gao, L. Huang, C. Zhang, An integrated biorefinery process: Stepwise extraction of fucoxanthin, eicosapentaenoic acid and chrysolaminarin from the same *Phaeodactylum tricornutum* biomass, *Algal Res.* 32 (2018) 193–200. <https://doi.org/10.1016/j.algal.2018.04.002>.
 - [27] A. Delbrut, P. Albina, T. Lapierre, R. Pradelles, E. Dubreucq, Fucoxanthin and Polyunsaturated Fatty Acids Co-Extraction by a Green Process, *Molecules.* 23 (2018) 874. <https://doi.org/10.3390/molecules23040874>.
 - [28] S.M. Kim, S.W. Kang, O.N. Kwon, D. Chung, C.H. Pan, Fucoxanthin as a major carotenoid in *Isochrysis aff. galbana*: Characterization of extraction for commercial application, *J. Korean Soc. Appl. Biol. Chem.* (2012).
<https://doi.org/10.1007/s13765-012-2108-3>.
 - [29] A.G. Pereira, P. Otero, J. Echave, A. Carreira-Casais, F. Chamorro, N. Collazo, A. Jaboui, C. Lourenço-Lopes, J. Simal-Gandara, M.A. Prieto, Xanthophylls from the Sea: Algae as Source of Bioactive Carotenoids, *Mar. Drugs.* 19 (2021) 188. <https://doi.org/10.3390/md19040188>.
 - [30] M. Bertrand, Carotenoid biosynthesis in diatoms, *Photosynth. Res.* 106 (2010) 89–102. <https://doi.org/10.1007/s11120-010-9589-x>.

- [31] M. Bae, M.-B. Kim, Y.-K. Park, J.-Y. Lee, Health benefits of fucoxanthin in the prevention of chronic diseases, *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids.* 1865 (2020) 158618. <https://doi.org/10.1016/j.bbalip.2020.158618>.
- [32] R. Vardanyan, V. Hruby, Hypolipidemic and Antihyperlipidemic Drugs, in: *Synth. Best-Seller Drugs*, Elsevier, 2016: pp. 285–315. <https://doi.org/10.1016/B978-0-12-411492-0.00020-1>.
- [33] L. Barsanti, P. Gualtieri, Is exploitation of microalgae economically and energetically sustainable?, *Algal Res.* 31 (2018) 107–115. <https://doi.org/10.1016/j.algal.2018.02.001>.
- [34] R. Sathasivam, R. Radhakrishnan, A. Hashem, E.F. Abd_Allah, Microalgae metabolites: A rich source for food and medicine, *Saudi J. Biol. Sci.* 26 (2019) 709–722. <https://doi.org/10.1016/j.sjbs.2017.11.003>.
- [35] F. Gao, M. Sá, I.T.D. Cabanelas, R.H. Wijffels, M.J. Barbosa, Improved fucoxanthin and docosahexaenoic acid productivities of a sorted self-settling *Tisochrysis lutea* phenotype at pilot scale, *Bioresour. Technol.* 325 (2021) 124725. <https://doi.org/10.1016/j.biortech.2021.124725>.
- [36] P. Steinrücken, S.K. Prestegard, J.H. de Vree, J.E. Storesund, B. Pree, S.A. Mjøs, S.R. Erga, Comparing EPA production and fatty acid profiles of three *Phaeodactylum tricornutum* strains under western Norwegian climate conditions, *Algal Res.* 30 (2018) 11–22. <https://doi.org/10.1016/j.algal.2017.12.001>.
- [37] S. Xia, B. Gao, J. Fu, J. Xiong, C. Zhang, Production of fucoxanthin, chrysolaminarin, and eicosapentaenoic acid by *Odontella aurita* under different nitrogen supply regimes, *J. Biosci. Bioeng.* 126 (2018) 723–729. <https://doi.org/10.1016/j.jbiosc.2018.06.002>.
- [38] S. Tachihana, N. Nagao, T. Katayama, M. Hirahara, F.M. Yusoff, S. Banerjee, M. Shariff, N. Kurosawa, T. Toda, K. Furuya, High Productivity of Eicosapentaenoic Acid and Fucoxanthin by a Marine Diatom *Chaetoceros gracilis* in a Semi-

- Continuous Culture, *Front. Bioeng. Biotechnol.* 8 (2020).
<https://doi.org/10.3389/fbioe.2020.602721>.
- [39] X. Mao, S.H.Y. Chen, X. Lu, J. Yu, B. Liu, High silicate concentration facilitates fucoxanthin and eicosapentaenoic acid (EPA) production under heterotrophic condition in the marine diatom *Nitzschia laevis*, *Algal Res.* 52 (2020) 102086.
<https://doi.org/10.1016/j.algal.2020.102086>.
- [40] Z.-Y. Wen, Y. Jiang, F. Chen, High cell density culture of the diatom *Nitzschia laevis* for eicosapentaenoic acid production: fed-batch development, *Process Biochem.* 37 (2002) 1447–1453. [https://doi.org/10.1016/S0032-9592\(02\)00034-1](https://doi.org/10.1016/S0032-9592(02)00034-1).
- [41] I. Tzovenis, N. De Pauw, P. Sorgeloos, Effect of different light regimes on the docosahexaenoic acid (DHA) content of *Isochrysis aff. galbana* (clone T-ISO), *Aquac. Int.* 5 (1997) 489–507. <https://doi.org/10.1023/A:1018349131522>.
- [42] G. Panis, J.R. Carreon, Commercial astaxanthin production derived by green alga *Haematococcus pluvialis*: A microalgae process model and a techno-economic assessment all through production line, *Algal Res.* 18 (2016) 175–190.
<https://doi.org/10.1016/j.algal.2016.06.007>.
- [43] D. Ippoliti, C. Gómez, M. del Mar Morales-Amaral, R. Pistocchi, J.M. Fernández-Sevilla, F.G. Acién, Modeling of photosynthesis and respiration rate for *Isochrysis galbana* (T-Iso) and its influence on the production of this strain, *Bioresour. Technol.* 203 (2016) 71–79.
<https://doi.org/10.1016/j.biortech.2015.12.050>.
- [44] H. Hu, L.-L. Ma, X.-F. Shen, J.-Y. Li, H.-F. Wang, R.J. Zeng, Effect of cultivation mode on the production of docosahexaenoic acid by *Tisochrysis lutea*, *AMB Express.* 8 (2018) 50. <https://doi.org/10.1186/s13568-018-0580-9>.
- [45] F. da Costa, F. Le Grand, C. Quéré, G. Bougaran, J.P. Cadoret, R. Robert, P. Soudant, Effects of growth phase and nitrogen limitation on biochemical composition of two strains of *Tisochrysis lutea*, *Algal Res.* 27 (2017) 177–189.

<https://doi.org/10.1016/j.algal.2017.09.003>.

- [46] H. Hu, J.Y. Li, X.R. Pan, F. Zhang, L.L. Ma, H.J. Wang, R.J. Zeng, Different DHA or EPA production responses to nutrient stress in the marine microalga *Tisochrysis lutea* and the freshwater microalga *Monodus subterraneus*, *Sci. Total Environ.* 656 (2019) 140–149. <https://doi.org/10.1016/j.scitotenv.2018.11.346>.
- [47] C.E. de Farias Silva, A. Bertucco, Bioethanol from microalgae and cyanobacteria: A review and technological outlook, *Process Biochem.* 51 (2016) 1833–1842. <https://doi.org/10.1016/j.procbio.2016.02.016>.
- [48] A. Hosseini, M. Jazini, M. Mahdиеh, K. Karimi, Efficient superantioxidant and biofuel production from microalga *Haematococcus pluvialis* via a biorefinery approach, *Bioresour. Technol.* 306 (2020) 123100. <https://doi.org/10.1016/j.biortech.2020.123100>.
- [49] R.R.L. Guillard, Culture of Phytoplankton for Feeding Marine Invertebrates, in: *Cult. Mar. Invertebr. Anim.*, 1975. https://doi.org/10.1007/978-1-4615-8714-9_3.
- [50] R.A. Andersen, J.A. Berges, P.J. Harrison, M.M. Watanabe, Recipes for Freshwater and Seawater Media, *Algal Cult. Tech.* (2005) 429–538. <https://doi.org/10.1016/b978-012088426-1/50027-5>.
- [51] Y. Collos, F. Mornet, A. Sciandra, N. Waser, A. Larson, P.J. Harrison, An optical method for the rapid measurement of micromolar concentrations of nitrate in marine phytoplankton cultures, *J. Appl. Phycol.* (1999). <https://doi.org/10.1023/A:1008046023487>.
- [52] Y.-K. Lee, W. Chen, H. Shen, D. Han, Y. Li, H.D.T. Jones, J.A. Timlin, Q. Hu, Basic Culturing and Analytical Measurement Techniques, in: *Handb. Microalgal Cult.*, John Wiley & Sons, Ltd, Oxford, UK, 2013: pp. 37–68. <https://doi.org/10.1002/9781118567166.ch3>.
- [53] X. Ma, J. Liu, B. Liu, T. Chen, B. Yang, F. Chen, Physiological and biochemical changes reveal stress-associated photosynthetic carbon partitioning into

- triacylglycerol in the oleaginous marine alga *Nannochloropsis oculata*, *Algal Res.* 16 (2016) 28–35. <https://doi.org/10.1016/j.algal.2016.03.005>.
- [54] G. Breuer, W.A.C. Evers, J.H. De Vree, D.M.M. Kleinegris, D.E. Martens, R.H. Wijffels, P.P. Lamers, Analysis of Fatty Acid Content and Composition in Microalgae, 5 (2013) 1–9. <https://doi.org/10.3791/50628>.
- [55] J. Jia, D. Han, H.G. Gerken, Y. Li, M. Sommerfeld, Q. Hu, J. Xu, Molecular mechanisms for photosynthetic carbon partitioning into storage neutral lipids in *Nannochloropsis oceanica* under nitrogen-depletion conditions, *Algal Res.* 7 (2015) 66–77. <https://doi.org/10.1016/j.algal.2014.11.005>.
- [56] M. DuBois, K.A. Gilles, J.K. Hamilton, P.A. Rebers, F. Smith, Colorimetric Method for Determination of Sugars and Related Substances, *Anal. Chem.* 28 (1956) 350–356. <https://doi.org/10.1021/ac60111a017>.
- [57] R. Sivaramakrishnan, A. Incharoensakdi, Utilization of microalgae feedstock for concomitant production of bioethanol and biodiesel, *Fuel.* 217 (2018) 458–466. <https://doi.org/10.1016/j.fuel.2017.12.119>.
- [58] A.W. Almutairi, Improvement of Chemical Composition of *Tisochrysis lutea* Grown Mixotrophically under Nitrogen Depletion towards Biodiesel Production, *Molecules.* 25 (2020) 4609. <https://doi.org/10.3390/molecules25204609>.
- [59] M. Garnier, G. Carrier, H. Rogniaux, E. Nicolau, G. Bougaran, B. Saint-Jean, J.P. Cadoret, Comparative proteomics reveals proteins impacted by nitrogen deprivation in wild-type and high lipid-accumulating mutant strains of *Tisochrysis lutea*, *J. Proteomics.* 105 (2014) 107–120. <https://doi.org/10.1016/j.jprot.2014.02.022>.
- [60] G. Carrier, M. Garnier, L. Le Cunff, G. Bougaran, I. Probert, C. De Vargas, E. Corre, J.-P. Cadoret, B. Saint-Jean, Comparative Transcriptome of Wild Type and Selected Strains of the Microalgae *Tisochrysis lutea* Provides Insights into the Genetic Basis, Lipid Metabolism and the Life Cycle, *PLoS One.* 9 (2014) e86889.

- <https://doi.org/10.1371/journal.pone.0086889>.
- [61] A. Charrier, J.-B. Bérard, G. Bougaran, G. Carrier, E. Lukomska, N. Schreiber, F. Fournier, A.F. Charrier, C. Rouxel, M. Garnier, J.-P. Cadoret, B. Saint-Jean, High-affinity nitrate/nitrite transporter genes (Nrt2) in *Tisochrysis lutea* : identification and expression analyses reveal some interesting specificities of Haptophyta microalgae, *Physiol. Plant.* 154 (2015) 572–590.
<https://doi.org/10.1111/ppl.12330>.
- [62] T. You, S.M. Barnett, Effect of light quality on production of extracellular polysaccharides and growth rate of *Porphyridium cruentum*, *Biochem. Eng. J.* 19 (2004) 251–258. <https://doi.org/10.1016/j.bej.2004.02.004>.
- [63] V.C. Liyanaarachchi, G.K.S.H. Nishshanka, R.G.M.M. Premaratne, T.U. Ariyadasa, P.H.V. Nimarshana, A. Malik, Astaxanthin accumulation in the green microalga *Haematococcus pluvialis*: Effect of initial phosphate concentration and stepwise/continuous light stress, *Biotechnol. Reports.* 28 (2020) e00538.
<https://doi.org/10.1016/j.btre.2020.e00538>.
- [64] Z. Sun, Y. Chen, X. Mao, J. Liu, Physiological and Biochemical Changes Reveal Differential Patterns of Docosahexaenoic Acid Partitioning in Two Marine Algal Strains of *Isochrysis*, *Mar. Drugs.* 15 (2017) 357.
<https://doi.org/10.3390/md15110357>.
- [65] F. Mus, J.-P. Toussaint, K.E. Cooksey, M.W. Fields, R. Gerlach, B.M. Peyton, R.P. Carlson, Physiological and molecular analysis of carbon source supplementation and pH stress-induced lipid accumulation in the marine diatom *Phaeodactylum tricornutum*, *Appl. Microbiol. Biotechnol.* 97 (2013) 3625–3642.
<https://doi.org/10.1007/s00253-013-4747-7>.
- [66] J.M. Roux, H. Lamotte, J.L. Achard, An Overview of Microalgae Lipid Extraction in a Biorefinery Framework, *Energy Procedia.* 112 (2017) 680–688.
<https://doi.org/10.1016/j.egypro.2017.03.1137>.

- [67] J. Liu, X. Mao, W. Zhou, M.T. Guarnieri, Simultaneous production of triacylglycerol and high-value carotenoids by the astaxanthin-producing oleaginous green microalga Chlorella zofingiensis, *Bioresour. Technol.* 214 (2016) 319–327. <https://doi.org/10.1016/j.biortech.2016.04.112>.
- [68] R. Gallego, C. Tardif, C. Parreira, T. Guerra, M.J. Alves, E. Ibáñez, M. Herrero, Simultaneous extraction and purification of fucoxanthin from *Tisochrysis lutea* microalgae using compressed fluids, *J. Sep. Sci.* 43 (2020) 1967–1977. <https://doi.org/10.1002/jssc.202000021>.
- [69] R. Gonçalves de Oliveira-Júnior, R. Grougnet, P.-E. Bodet, A. Bonnet, E. Nicolau, A. Jebali, J. Rumin, L. Picot, Updated pigment composition of *Tisochrysis lutea* and purification of fucoxanthin using centrifugal partition chromatography coupled to flash chromatography for the chemosensitization of melanoma cells, *Algal Res.* 51 (2020) 102035. <https://doi.org/10.1016/j.algal.2020.102035>.
- [70] B. Aslanbay Guler, I. Deniz, Z. Demirel, O. Yesil-Celiktas, E. Imamoglu, A novel subcritical fucoxanthin extraction with a biorefinery approach, *Biochem. Eng. J.* 153 (2020) 107403. <https://doi.org/10.1016/j.bej.2019.107403>.
- [71] S. Garg, S. Afzal, A. Elwakeel, D. Sharma, N. Radhakrishnan, J.K. Dhanjal, D. Sundar, S.C. Kaul, R. Wadhwa, Marine Carotenoid Fucoxanthin Possesses Anti-Metastasis Activity: Molecular Evidence, *Mar. Drugs.* 17 (2019) 338. <https://doi.org/10.3390/md17060338>.
- [72] S. Deshmukh, R. Kumar, K. Bala, Microalgae biodiesel: A review on oil extraction, fatty acid composition, properties and effect on engine performance and emissions, *Fuel Process. Technol.* 191 (2019) 232–247. <https://doi.org/10.1016/j.fuproc.2019.03.013>.
- [73] I. Pancha, K. Chokshi, B. George, T. Ghosh, C. Paliwal, R. Maurya, S. Mishra, Nitrogen stress triggered biochemical and morphological changes in the

- microalgae *Scenedesmus* sp. CCNM 1077, *Bioresour. Technol.* 156 (2014) 146–154. <https://doi.org/10.1016/j.biortech.2014.01.025>.
- [74] D. Cheng, D. Li, Y. Yuan, L. Zhou, X. Li, T. Wu, L. Wang, Q. Zhao, W. Wei, Y. Sun, Improving carbohydrate and starch accumulation in *Chlorella* sp. AE10 by a novel two-stage process with cell dilution, *Biotechnol. Biofuels.* 10 (2017) 75. <https://doi.org/10.1186/s13068-017-0753-9>.
- [75] L.E. Walls, S.B. Velasquez-Orta, E. Romero-Frasca, P. Leary, I. Yáñez Noguez, M.T. Orta Ledesma, Non-sterile heterotrophic cultivation of native wastewater yeast and microalgae for integrated municipal wastewater treatment and bioethanol production, *Biochem. Eng. J.* 151 (2019) 107319. <https://doi.org/10.1016/j.bej.2019.107319>.
- [76] P. Choudhary, P.P. Assemany, F. Naaz, A. Bhattacharya, J. de S. Castro, E. de A. do C. Couto, M.L. Calijuri, K.K. Pant, A. Malik, A review of biochemical and thermochemical energy conversion routes of wastewater grown algal biomass, *Sci. Total Environ.* 726 (2020) 137961. <https://doi.org/10.1016/j.scitotenv.2020.137961>.