

A PRELIMINARY STUDY  
OF THE  
SPATIAL VARIATION OF RAINFALL  
AND ITS EFFECT ON  
WATER MANAGEMENT IN THE DRY ZONE

by

P.S. RAVINDRA

B.Sc.(Eng.) Hons., M.I.E. (SL), MICE, C Eng.

A dissertation submitted in partial fulfilment  
of the requirements for the degree of  
Master of Engineering

Department of Civil Engineering,  
University of Moratuwa,  
Sri Lanka.

March 1985

54102

This dissertation has not been previously presented in whole or part, to any University or Institution for a higher degree.

A handwritten signature in black ink, appearing to be 'P.S. Ravindra', written over a horizontal dotted line.

P.S. Ravindra

March 1985.

ABSTRACT

Incorporating effective rainfall in irrigation water is an important task in water management. Understanding the climatic conditions is necessary to plan a raingauge network. The occurrence of thunderstorms is a main feature in the inter-monsoonal periods. These convective storms are often highly localized and usually tend to be less than 8 km in width. Thus a fairly dense network of raingauges is necessary to get an accurate representation of the rainfall during these periods. However, when the rainfall is widespread, the maintenance of such a dense raingauge network is not essential.

Paddy production in Sri Lanka is more influenced by rainfall in the Maha (October to February) season than in the Yala (May to August), as there is hardly any rain in the Yala season in most of the paddy growing areas. For the Maha crop, land preparation usually starts in October. About 28 percent of the annual rainfall occurs in October and November (second inter-monsoonal period) and is mostly of the convective type. With a proper understanding of the areal rainfall distribution during this period, conserving irrigation water is possible.

The use of isohyetal patterns, correlation coefficient vs. distance diagrams and correlation linkage diagrams could be used to assess an optimum spacing between the raingauges.

Using a triangular storm model, it can be shown that negative correlation values between pairs of raingauges, are due to smaller storm cells. Further, it can be shown when the storm width is approximately three times the width of the gauged area then no negative correlation can occur between the raingauges.

A square grid of raingauges at about 2 km intervals is necessary during the period of the convective storms while a 14 km grid is required for the widespread monsoonal storm period.

This study indicates an area where more research work is required for better water management and it is possible to achieve this within the normal level of management and would also yield considerable economic benefits.

ACKNOWLEDGEMENTS

Thanks are due to several persons who helped make this study a success; Dr. Sunil Wickremasuriya the course co-ordinator for generously contributing his time and effort in guiding me throughout the project, Mr. N. Kumarasamy, Senior Deputy Director (Water Management), Irrigation Department for his personal guidance and for making available the resources of the department, and Mr. J.T. Arulanantham, Director of Meteorology for providing me with the relevant literature and for his comments and encouragement.

Thanks are also due to Mr. U.S. Weerasekera, Mr. P.P. Gnanapala, Mr. R.P. Lokuratna and Mr. G.J. Wickramage who gave their unstinted support to this study and contributed variously throughout its duration.

List of Symbols and Abbreviations

$r$	-	correlation coefficient
$\Sigma$	-	summation
$X, Y$	-	precipitation observation series
$N$	-	number of observations
$\rho (D)$	-	inter-station correlation function
$D$	-	distance between raingauges
$B$	-	storm cell diameter
$G_i$	-	measuring gauge $i$
$g_i(t)$	-	rainfall amount produced by storm $t$
$S_c$	-	storm centre
$h_a$	-	rainfall amount produced at point 'a' of the storm
$M_{in}$	-	minimum value
$Y_i$	-	gauge axis
$Z_i$	-	area axis

TABLE OF CONTENTS

ABSTRACT	I
ACKNOWLEDGEMENTS	III
LIST OF SYMBOLS AND ABBREVIATIONS	IV
<u>1. INTRODUCTION</u>	1-4
1.1 General	1
1.2 Objective of Study	2
1.3 Method of Study	4
<u>2. RAINFALL PATTERN IN SRI LANKA</u>	5-14
2.1 Mean Annual Rainfall of Sri Lanka	5
2.2 Weather Pattern	8
<u>3. SPATIAL VARIATION IN RAINFALL AND     ITS EFFECT ON PADDY PRODUCTION</u>	15-21
3.1 Land Preparation Requirement	15
3.2 Crop Water Requirement	17
3.3 Effective Rainfall	18
3.4 Effect of Spatial Variation of Rainfall on Rice Yield	19
<u>4. EFFICIENCY OF THE DENSITY OF RAINGAUGE NETWORKS</u>	22-37
4.1 Design of Raingauge Network	22
4.2 Isohyetal Maps	25
4.3 Distance vs. Correlation Coefficient Diagrams between the Raingauge Stations	26
4.4 Linear Distance Model	30
4.5 Correlation Linkage Diagrams	35

<u>5.</u>	<u>MATHEMATICAL MODEL FOR A STORM</u>	38-43
5.1	General	38
5.2	Model for a Storm using the Triangular Distribution	39
<u>6.</u>	<u>DISCUSSION OF RESULTS</u>	44-47
6.1	General	44
6.2	The Raingauge Distance vs. Correlation Coefficient Graphs	44
6.3	Fitting of Regression Curve	45
6.4	Correlation Linkage Diagrams	46
6.5	Isohyetal Patterns	46
<u>7.</u>	<u>CONCLUSIONS</u>	48-50
7.1	Conclusions	48
7.2	Future Study	49
<u>8.</u>	<u>REFERENCES</u>	51
<u>9.</u>	<u>PLATES</u>	53-57
<u>10.</u>	<u>APPENDICES</u>	58-146
10.1	Appendix 1	58
10.2	Appendix 2	109
10.3	Appendix 3	115